

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Atty. Dkt. No. 072827-1905

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently amended) A method of treating a patient for depression comprising ~~the step of administering to said patient an effective amount of~~ selecting a compound having a NMDA IC₅₀ of about 50 nM to about 1 μM as measured in the NMDA assay and a serotonin reuptake IC₅₀ of less than or equal to about 100 nm as measured in the serotonin reuptake inhibition assay; and
administering to said patient an effective amount of said compound.
2. (Original) The method of claim 1, wherein said compound has an NMDA receptor IC₅₀ of 50 nM to 1 μM and a SSRI IC₅₀ less than 100 nM.
3. (Currently amended) A method of treating a patient for depression comprising ~~the step of administering to said patient an effective amount of~~ selecting a compound a compound having a NMDA IC₅₀ of about 50 nM to about 1 μM as measured in the NMDA assay and a serotonin reuptake IC₅₀ of less than or equal to about 100 nm as measured in the serotonin reuptake inhibition assay, wherein said compound has having the chemical structure:



wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, and -O-acyl;

Ar¹ and Ar² are each independently selected from the group consisting of phenyl, naphthyl, thiofuranyl, tetrahydronaphthyl, furanyl, tetrahydrofuranyl, pyridyl, quinoliny,

Atty. Dkt. No. 072827-1905

cyclopentyl, tetrahydroquinoliny, tetrahydroisoquinoliny, cyclohexyl, cycloheptyl, and cyclopentyl;

each R^1 is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;

each R^2 is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R^2 's together are imino;

each R^3 is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl; and

each m is independently an integer from 0 to 5;

provided that if both R_3 's are $-CH_3$, then both X_m 's are not 3-F, 4-F, 3- CF_3 , 4-Cl, and if both R_3 's are $-CH_3$ and one X_m is 4-F then the other X_m is not 4-Cl; further provided that if one R_3 is -H and the other R_3 is $-CH_3$ then both X_m 's are not 4-Cl, and if one R_3 is -H and the other R_3 is $-CH_3$ then at least one m is 1;

or a pharmaceutically acceptable salt thereof.

4. (Previously presented) The method of claim 3 wherein for said compound each X is independently either -F, -Cl, $-OCF_3$ or $-CF_3$;

each R^1 is -H;

each R^2 is -H;


one R^3 is -H, and the other R^3 is either -H or $-CH_3$; and

each m is 1.

5. (Currently amended) A method of treating a patient for depression comprising administering to said patient an effective amount of a compound having the chemical structure

~~The method of claim 3 wherein said compound has the chemical structure:~~

Atty. Dkt. No. 072827-1905



wherein X^1 is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, —OCF₃, -O-alkyl, or -O-acyl;


X^2 is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, —OCF₃, -O-alkyl, or -O-acyl; and

R^3 is either -H or -CH₃;

or a pharmaceutically acceptable salt thereof.

6. (Original) The method of claim 5, wherein X^1 is -F, -Cl, -OCF₃ or -CF₃; and X^2 is either 2-OCH₃, 2-CH₃, 3-F, 3-CF₃, or 4-CF₃.

7. (Withdrawn) A method of treating a patient for depression comprising the step of administering to said patient an effective amount of a compound having the chemical structure:



wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, —OCF₃, -O-alkyl, and -O-acyl;

W is selected from the group consisting of -CH₂, -O-, and -S-;

Ar^1 and Ar^2 are each independently selected from the group consisting of phenyl, naphthyl, thiofuranyl, tetrahydronaphthyl, furanyl, tetrahydrofuranyl, pyridyl, quinoliny,

Atty. Dkt. No. 072827-1905

isoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl cyclohexyl, cycloheptyl, and cyclopentyl;

each R^1 is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;

each R^2 is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R^2 's together are imino;

each R^3 is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl; and

m is 0 to 5;

or a pharmaceutically acceptable salt thereof.

8. (Withdrawn) The method of claim 7, wherein for said compound each X is independently either -F, -Cl, -OCF₃ or -CF₃;

Ar¹ and Ar² are each independently phenyl or naphthyl;


each R^1 is -H;

each R^2 is -H;

one R^3 is -H, and the other R^3 is either -H or -CH₃;

each m is 0 or 1.

9. (Withdrawn) The method of claim 7, wherein said compound has the chemical structure:




wherein X¹ is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl;

Atty. Dkt. No. 072827-1905

X^2 is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃,
 -O-alkyl, or -O-acyl; and
 R^3 is either -H or -CH₃;
 or a pharmaceutically acceptable salt thereof.

10. (Withdrawn) The method of claim 9 wherein X^1 is either -F, -Cl, -OCF₃ or -CF₃; and
 X^2 is either 2-OCH₃, 2-CH₃, 3-F, 3-CF₃, or 4-CF₃.

11. (Withdrawn) A method of treating a patient for depression comprising the step of
 administering to said patient an effective amount of a compound having the chemical structure:



wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃,

-O-alkyl, and -O-acyl;

each R¹ is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;

each R² is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R²'s together are imino;

each R³ is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl;

Z is either -CH₂CH₂-, -CH₂CH(CH₃)-, -CH=CH-, -O-CH₂-, -S-CH₂-, -CH₂-, -O-, or -S-;
 and

Atty. Dkt. No. 072827-1905

each n is independently 1 to 4; or a pharmaceutically acceptable salt thereof.

12. (Withdrawn) The compound of claim 11, wherein each X is independently either -F, -Cl, -OCF₃ or -CF₃;


each R¹ is -H;

each R² is -H;

one R³ is -H, and the other R³ is either -H or -CH₃; and

each n is 1.

13. (Withdrawn) The method of claim 11, wherein said compound has the chemical structure:



wherein X¹ is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl;

X² is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃,

-O-alkyl, or -O-acyl; and


R³ is either -H or -CH₃;

or a pharmaceutically acceptable salt thereof.

14. (Withdrawn) The method of claim 13 wherein X¹ is -F, -Cl, -OCF₃ or -CF₃; and X² is either -F, -Cl, -OCH₃, -CH₃, -OCF₃ or -CF₃.

Atty. Dkt. No. 072827-1905

15. (Withdrawn) A method of treating a patient for depression comprising the step of administering to said patient an effective amount of a compound having the chemical structure:



wherein each X is independently selected from the group consisting of -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃,

-O-alkyl, and -O-acyl; preferably, each X is independently either -F, -Cl, -OCF₃ or -CF₃;

Ar¹ and Ar² are each independently selected from the group consisting of phenyl, naphthyl, thiofuranyl, tetrahydronaphthyl, furanyl, tetrahydrofuranyl, pyridyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, cyclohexyl, cycloheptyl, and cyclopentyl; preferably Ar¹ and Ar² are independently naphthyl or phenyl; more preferably at least one of Ar¹ and Ar² is phenyl; and more preferably, both Ar¹ and Ar² are phenyl;

Y is either -CH₂-, -O-, or -S-;

each R¹ is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl; preferably, each R¹ is -H;

each R² is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R²s together are imino; preferably each R² is -H;


each R³ is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl; preferably, each R³ is independently either -H or -CH₃; more preferably one R³ is

-H, and the other R³ is either -H or -CH₃; and

each m is independently an integer from 0 to 5; and preferably, each m is independently 0 or 1.

16. (Withdrawn) The method of claim 15, wherein said compound has the chemical structure; Structure VIII

Atty. Dkt. No. 072827-1905



wherein X^1 is independently selected from the group consisting of $-H$, $-Br$, $-Cl$, $-F$, $-I$, $-CF_3$, alkyl, $-OH$,

$-OCF_3$, $-O$ -alkyl, or $-O$ -acyl; preferably, X^1 is either $-F$, $-Cl$, $-OCF_3$ and $-CF_3$;

X^2 is either $-Br$, $-Cl$, $-F$, $-I$, $-CF_3$, alkyl, $-OH$, $-OCF_3$,

$-O$ -alkyl, or $-O$ -acyl; preferably, X^2 is independently either $-F$, $-Cl$, $-OCH_3$, $-CH_3$, $-OCF_3$ or $-CF_3$;














more preferably, X^2 is either 2- OCH_3 , 2- CH_3 , 3- F , 3- CF_3 , or 4- CF_3 ; and

R^3 is either $-H$ or CH_3 ;

or a pharmaceutically acceptable salt thereof.

17. (Withdrawn) A compound having the chemical structure;














Atty. Dkt. No. 072827-1905



or a pharmaceutically acceptable salt thereof.

Atty. Dkt. No. 072827-1905


18. (Withdrawn) A method of treating a patient for depression comprising the step of administering to said patient an effective amount of a compound having the chemical structure:



Atty. Dkt. No. 072827-1905

or a pharmaceutically acceptable salt thereof.

19. (Previously presented) The method of claim 3 wherein said compound has the chemical structure:



wherein

X¹ is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl;

X² is either -Br, -Cl, -F, -I, -CF₃, alkyl, -OH, -OCF₃, -O-alkyl, or -O-acyl;

each R¹ is independently selected from the group consisting of -H, alkyl, hydroxyalkyl, -OH, -O-alkyl, and -O-acyl;

each R² is independently selected from the group consisting of -H, alkyl, and hydroxyalkyl, or both R²'s together are imino

each R³ is independently selected from the group consisting of -H, alkyl, 2-hydroxyethyl, and alkylphenyl;

or a pharmaceutically acceptable salt thereof.

20. (Previously presented) The method of claim 19, wherein

each X is independently either -F, -Cl, -OCF₃ or -CF₃;

each R¹ is -H;

each R² is -H; and

one R³ is -H, and the other R³ is either -H or -CH₃.

Atty. Dkt. No. 072827-1905

- C1
21. (New) The method of claim 5, wherein X^1 and X^2 are F, and R^3 is -H.
21. (New) The method of claim 21, wherein X^2 is at the 3-position.
23. (New) The method of claim 5, wherein X^1 and X^2 are F, and R^3 is -CH₃.
24. (New) The method of claim 23, wherein X^2 is at the 3-position.
-